



# The 1<sup>st</sup> International Neuroinflammation Congress and 1<sup>st</sup> Student Festival of Neuroscience

Shefa Neuroscience Research Center, Tehran, Iran, 11-13 April, 2017

*The Neuroscience Journal of Shefaye Khatam*

Volume 5, No. 2, Suppl 2

## Poster Presentation

### The Role of Neuroinflammation in Post-Traumatic Stress Disorder (PTSD)

Farideh Baghishani<sup>1\*</sup>, Sajad Sahab Negah<sup>2,3</sup>, Ehsan Baghishani<sup>4</sup>

<sup>1</sup>Department of Anatomy and Cell Biology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>2</sup>Department of Neuroscience, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>3</sup>Shefa Neuroscience Research Center, Khatam Alanbia Hospital, Tehran, Iran

<sup>4</sup>Islamic Azad University, Mashhad Branch, Mashhad, Iran

**Published: 11 April, 2017**

#### Abstract

Post-traumatic stress disorder (PTSD) is an anxiety disorder that can typically occur after traumatic events in which the individual felt severe helplessness or horror. PTSD has a lifetime prevalence of 5–8% and women show higher prevalence than in compared with men. Neuroinflammation is associated to anxiety and related disorders such as PTSD. It is an early, specialized immune reaction following tissue damage and/or pathogen invasion in the central nervous system (CNS). Regarding the importance of anxiety disorders especially PTSD and their effect on patients' personal and social life, considering of the neuroinflammation in PTSD can be used for new treatment. Several studies have been reported that there is a relationship between neuroinflammation and PTSD. Current epidemiological evidence shows that increased expression of pro-inflammatory cytokines such as interleukin-1 $\beta$ , interleukin-6, and tumor necrosis factor- $\alpha$ , and decreased anti-inflammatory factors have a key role in creating PTSD. Also, neuroinflammation induced by inhibition of the NADPH oxidase (NOX2) can lead to trigger the PTSD symptoms. Various inflammatory markers such as Cortisol, C-reactive protein (CRP), Th1 cytokines, and Th2 Cytokines are associated with PTSD. Furthermore, microglial activation may also initiate inflammation and can be involved in PTSD. Thus, scrutinized research about PTSD and understanding of creating mechanisms of this disorder is essential and can provide grounds for its treatment.

**Keywords:** Neuroinflammation, Post-traumatic stress disorder, Central nervous system

**\*Corresponding Author:** Farideh Baghishani

**E-mail:** baghishanif2@mums.ac.ir