



# The 2<sup>nd</sup> International Neuroinflammation Congress and 2<sup>nd</sup> Student Festival of Neuroscience

Shefa Neuroscience Research Center, Tehran, Iran, 17-19 April, 2018

*The Neuroscience Journal of Shefaye Khatam*

Volume 6, No. 2, Suppl 1

## Poster Presentation

### Dextran Curcumin Promotes Novel Object Recognition Memory in Rats after Ischemic Stroke

Homa Talebi<sup>1</sup>, Akbar Hajizadeh Moghaddam<sup>1</sup>, Mahbobe Zare<sup>2\*</sup>, Sedigheh Khanjani Jelodar<sup>3</sup>

<sup>1</sup>Faculty of Basic Sciences, University of Mazandaran, Babolsar, Iran

<sup>2</sup>Faculty of Herbs, Amol University of Special Modern Technologies, Amol, Iran

<sup>3</sup>Faculty of Biology, Shahid Beheshti University, Tehran, Iran

*Published: 17 April, 2018*

#### Abstract

Ischemic stroke causes the depletion of energy and induces excitotoxicity and neuroinflammation in the brain that results from thrombotic blockage. Cerebral ischemia leads to many types of memory loss, including impairment of working, spatial and object recognition memories. Curcumin shows strong anti-oxidoinflammatory activities but its therapeutic use is limited by its low solubility in water and corresponding poor intestinal absorption. So, in this study curcumin was used in conjugate with dextran as polymeric carriers in novel drug delivery system. The purpose of this study was to determine the effect of dextran-curcumin on memory impairment induced by global ischemia. In this study 35 rats divided into 5 groups. Pre-treatment and positive control groups, were treated with curcumin and dextran-curcumin (15mg/kg - orally) for 30 days and the vehicle and disease groups received distilled water. For induction of ischemic stroke model, rats were anesthetized and both right and left carotid arteries were selected and clamped for 5 min by vascular clamps (time of ischemia). Thereafter the vascular clamps were removed for the next 10 min (time of reperfusion), and both carotid arteries were clamped again for 5 min. Finally, the vascular clamps were removed and blood circulation was returned in both carotid arteries, 48 hours after induction of model, Novel Object Recognition test was used to determine memory impairment in all rats. Our study indicated that memory impairment increased in ischemic group and dextran curcumin has memory-improving effects after global ischemic stroke ( $p < 0.01$ ): Dextran-curcumin has memory-improving capacity better than curcumin in lower doses.

**Keywords:** Dextran-Curcumin, Memory, Ischemic Stroke

**\*Corresponding Author:** Mahbobe Zare

**E-mail:** mahboobeh.zare93@gmail.com