

# The 2<sup>nd</sup> International Neuroinflammation Congress and 2<sup>nd</sup> Student Festival of Neuroscience



Shefa Neuroscience Research Center, Tehran, Iran, 17-19 April, 2018

*The Neuroscience Journal of Shefaye Khatam*

Volume 6, No. 2, Suppl 1

## Poster Presentation

### Relationship Among Dimensions of Roy Adaptation Model, General Health and Satisfaction with Life in Parkinson Disease Patients

Monir Nobahar<sup>1\*</sup>, Hassan Babamohamadi<sup>2</sup>, Mohaddeseh Saffari<sup>2</sup>, Seyed Afshin Samaei<sup>3</sup>,  
Majid Mirmohammadkhani<sup>4</sup>

<sup>1</sup>Nursing Care Research Center, and Social Determinants of Health Research Center, Faculty of Nursing and Midwifery, Semnan University of Medical Sciences, Semnan, Iran

<sup>2</sup>Nursing Care Research Center, Faculty of Nursing and Midwifery, Semnan University of Medical Sciences, Semnan, Iran

<sup>3</sup>Rehabilitation Research Center, Neurology Department, Kowsar Hospital, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran

<sup>4</sup>Social Determinants of Health Research Center, Community Medicine Department, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran

*Published: 17 April, 2018*

#### Abstract

Parkinson disease (PD) is a chronic, progressive and debilitating disease. This study aimed to determine the relationship between the dimensions of Roy Adaptation Model (RAM) and General Health (GH) and Satisfaction with Life (SWL) in PD patients. This was a cross-sectional study. Samples were selected from the PD patients in 2016. The data was gathered by demographic, RAM, GH and SWL Scale questionnaires. In PD patients, total adaptation score was  $21.26 \pm 3.24$ . The dimensions of adaptation, the scores of physiologic mode, self-concept, role function and interdependence were  $74.7 \pm 9.33$ ,  $32.37 \pm 6.04$ ,  $24.30 \pm 5.00$  and  $20.93 \pm 2.46$ , respectively. There was significant relationship between education and physiologic mode ( $P=0.048$ ) and there were significant relationships between PD experience, employment and gender and total adaptation score ( $P=0.002$ ,  $P=0.007$ ,  $P=0.006$ , respectively). Mean and standard deviation of total general health score was  $30.97 \pm 5.03$  and the means and standard deviations of its dimensions, including somatic symptoms, anxiety and insomnia, social dysfunction and depression were  $10.26 \pm 2.71$ ,  $9.73 \pm 2.77$ ,  $5.13 \pm 2.23$  and  $5.83 \pm 1.91$ , respectively. There were significant relationships between social dysfunction and physiologic mode and self-concept ( $P=0.024$  and  $P=0.012$ , respectively). The mean and standard deviation of satisfaction with life score was  $21.47 \pm 3.57$ . Satisfaction with life had no relationship with dimensions of RAM and general health. RAM, GH and SWL make the development of high quality care planning possible by assessing the health status of PD patients. This leads to engagement of the patients in self-care, better adaptation, improved quality of life and also helps the nurses to make medical decisions.

**Keywords:** Roy Adaptation Model, General Health, Satisfaction with Life, Parkinson

**\*Corresponding Author:** Monir Nobahar

**Email:** nobahar43@yahoo.com