

The 2nd International Neuroinflammation Congress and 2nd Student Festival of Neuroscience



Shefa Neuroscience Research Center, Tehran, Iran, 17-19 April, 2018

The Neuroscience Journal of Shefaye Khatam

Volume 6, No. 2, Suppl 1

Poster Presentation

Inflammation or Neurodegeneration: Which one has Remarkable Role in Multiple Sclerosis?

Fahimeh Nourbakhsh^{1*}, Samira Barangi¹, Vajiheh Nourbakhsh², Afsaneh Yazdanpanah³, Elaheh Tajbakhsh⁴

¹Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

²Hazrat Fatemeh Zahra (SA) Hospital, Isfahan, Iran

³School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Department of Microbiology, Faculty of Basic Sciences, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

Published: 17 April, 2018

Abstract

Multiple Sclerosis (MS) is a complex disease resulting from the occurrence of intermingled episodes of neuroinflammation and degeneration. However, this concept has recently challenged by several observations suggesting that in this disease neurodegeneration might occur independently of inflammation. The evidence that active neurodegeneration in MS is invariably associated with inflammation is provided. The attack of myelin starts inflammatory processes, which triggers other immune cells and the release of soluble factors like cytokines and antibodies. Further breakdown of the blood–brain barrier in turn causes a number of other damaging effects such as swelling, activation of macrophages, more activation of cytokines and other destructive proteins. Inflammation can potentially reduce transmission of information between neurons in at least three ways. The soluble factors released might stop neurotransmission by intact neurons. These factors could lead to or enhance the loss of myelin, or they may cause the axon to break down completely. Neuroinflammation could also found in deep gray matter with pathological and clinical relevance. Therefore, control of inflammation with anti-inflammatory therapies must take into account as one of the main purposes of MS treatment parallel with other immunomodulatory and immunosuppressive treatments. Future therapeutic options for this disease discussed based on recent knowledge of the mechanisms of inflammation and neurodegeneration.

Keywords: Inflammation, Neurodegeneration, Multiple Sclerosis

***Corresponding Author:** Fahimeh Nourbakhsh

E-mail: nourbakhshf951@mums.ac.ir