

The 2nd International Neuroinflammation Congress and 2nd Student Festival of Neuroscience



Shefa Neuroscience Research Center, Tehran, Iran, 17-19 April, 2018

The Neuroscience Journal of Shefaye Khatam

Volume 6, No. 2, Suppl 1

Poster Presentation

Low-Intensity Aerobic Training for Along with Blood Flow Restriction on Amount of Protein BDNF in Soleus and EDL Muscles as well as the Sciatic Nerve in Aged Male Rats

Mohammad-Ali Bahreini Pour^{1*}, Fariborz Hovanloo², Siyavash Joukar^{3,4,5,6}, Hamid Najafipour^{5,6}

¹Department of Physical Education, Faculty of Shahid Chamran, Technical and Vocational University (YVU), Tehran, Iran

²Physical Education and Sports Science College, Shahid Beheshti University, Tehran, Iran

³Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran

⁴Physiology Research Center, Institute of Basic and Clinical Physiology Sciences, Kerman University of Medical Sciences, Kerman, Iran

⁵Cardiovascular Research Center, Institute of Basic and Clinical Physiology Sciences, Kerman University of Medical Science, Kerman, Iran

⁶Department of Physiology and Pharmacology, Kerman University of Medical Sciences, Kerman, Iran

Published: 17 April, 2018

Abstract

Neurotrophins are a group of nerve growth factors with their protein structure. These proteins play an important role in the growth and metabolism of many cells and maintain nerve, muscle, synaptic potentials, and also helps reduce depression and apoptosis. The aim of this study was to evaluate the effect of 10 weeks low-intensity aerobic training and limited blood flow to the amount of protein neurotrophic factor derived from the brain (BDNF) in the soleus, extensor long fingers muscles (EDL) as well as sciatic nerve in rats the elderly. 60 elderly male Wistar rats (23-24 months) were chosen which their weigh were 355 to 481 grams. They were divided into 6 groups randomly, blood flow restriction (BFR), exercise with blood flow restriction (BFR + Ex), sham (Sh), sham with exercise (Sham + Ex), control (Ctl) and exercise (Ex). They were trained with a low aerobic exercise (15 m / min) for 1 hour, 5 days a week for 10 weeks then were sacrificed 48 hours after the last training session. The samples of muscle and the sciatic nerve were separated immediately and they were put in a solid nitrogen. and Then, they were preserved at -80 ° C. The protein samples were detected by Western Blotting method. The statistical analysis was conducted with SPSS software version 18. Data normal distribution determined by Shapiro – Wilk test. Comparisons were performed among different groups by one-way ANOVA and post hoc Tukey's test. P values less than 0.05 were considered as statistically significant. The results have shown that BDNF protein just decreased in the EDL muscle for BFR + Ex group compared with Ctl and Sh groups (P<0.05) significantly. In addition, in soleus muscle BFR + Ex group decreased in compare with all groups (groups Ctl, Sh and BFR P< 0.001 and Ex and Sh + Ex (p<0.05) significantly. moreover, Ex, Sh + Ex groups in comparison with Ctl and Sh have decreased significantly (P< 0.001). On the other hand, the sciatic nerve of BFR + Ex group, in comparison with all groups, increased significant (P< 0.001). This study has indicated that this type of exercise can effect on the BDNF protein in slow and fast witch muscles. Furthermore, BFR+Ex, as a training method, can influence on nerve and muscle positively.

Keywords: Endurance Training, Neurotrophic Factors, Reduced Blood Flow- Aging

***Corresponding Author:** Mohammad Ali Bahreini Pour

Email: alibahreini1974@gmail.com