ثفار فق

The 1st International Neuroinflammation Congress and 1st Student Festival of Neurosience

Shefa Neuroscience Research Center, Tehran, Iran, 11-13 April, 2017

The Neuroscience Journal of Shefaye Khatam

Volume 5, No. 2, Suppl 2

Oral Presentation

Childhood Anti-NMDA Receptor Encephalitis

Farah Ashrafzadeh*

Mashhad University of Medical Sciences, Mashhad, Iran

Published: 11 April, 2017

Abstract

Anti N-methyl-D-aspartate receptor (NMDR) encephalitis has been recognized as the most frequent autoimmune encephalitis in children after acute demyelinating encephalomyelitis (ADEM). However due to the variable the variable clinical presentation, the paucity of specific finding on standard laboratory and radiological investigation remains under recognized. First discribed in 2005, most commonly affected are children and young adults; 80% patients are females. It has characteristic evolution in several stages, in prodromal phase appears fever, malaise, nausea, vomiting, diarrhea, or upper respiratory tract symptoms in about 70%, then gradually manifest psychotic symptoms (delusions, hallucination and mania) that in children can be present as temper tantrum, hyperactivity, irritability, neurologic presentation as seizure, status epilepticus and dystonia. NMDA receptors are ligand cation channels involved in synaptic glutaminergic transmission that plays a key role in functions such as memory, learning, behavior and cognition. Anti-NMDA encephalitis is associated antibodies against the NRII subunit of NMDA receptors. MRI changes are non-specific; EEG changes are slowing and background activity; CSF abnormalities are common, mild lymphocytic pleocytosis, elevated protein and positive oligoclonal bands in 60% of patients. Definitive diagnosis is based on demonstration of anti-NMDA antibodies in CSF or serum. Management include symptomatic therapy, definitive immunotherapy, and tumor surveillance. Early diagnosis and aggressive immunetherapy are important. In this presentation, I introduce a patient with anti-NMDA encephalitis.

Keywords: Childhood, Patients, Management

*Corresponding Author: Farah Ashrafzadeh

E-mail: ashrafzadehf@mums.ac.ir

