



# The 1<sup>st</sup> International Neuroinflammation Congress and 1<sup>st</sup> Student Festival of Neuroscience

Shefa Neuroscience Research Center, Tehran, Iran, 11-13 April, 2017

*The Neuroscience Journal of Shefaye Khatam*

Volume 5, No. 2, Suppl 2

## Oral Presentation

### Treatment of Traumatic Brain Injury in Adult Rats with Injection of Human Epileptic Neural Stem Cells and Nano-Scaffold

Sajad Sahab Negah<sup>1,2</sup>, Ali Jahanbazi Jahan-Abad<sup>2,3</sup>, Ali Gorji<sup>1,2,4\*</sup>

<sup>1</sup>Department of Neuroscience, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>2</sup>Shefa Neuroscience Research Center, Khatam Alanbia Hospital, Tehran, Iran

<sup>3</sup>Department of Clinical Biochemistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>4</sup>Epilepsy Research Center, Department of Neurology, and Department of Neurosurgery, Westfälische Wilhelms-Universität Münster, Münster, Germany

**Published: 11 April, 2017**

#### Abstract

Traumatic brain injury (TBI) is described by a disruption in the normal function of the brain due to an injury following a trauma, which can potentially cause severe physical, cognitive, and emotional impairment. The use of human stem cells and self-assembling peptide scaffolds suggest huge potential for application in the treatment of TBI. In the present study, we surveyed the beneficial effects of human adipose-derived stem cells (hADSCs), human epileptic neural stem cells (hENSCs), and PuraMatrix hydrogel (PM) in an acute brain injury model of mild TBI using 3 months rats. hADSCs and hENSCs were transplanted with or without scaffolds into rats brain and the electroencephalography were compared. Modified neurologic severity score (mNSS) tests were performed to measure behavioral outcomes. PM scaffold increased the retention of hENSCs in the lesion site and limited its distribution at the transplanted region. Significantly more hENSCs were detected in the brain when transplanted with PM scaffold. The results showed PM scaffold also efficiently improved cell survival *in vivo*, resulting in better neural functional recovery. We hypothesized that PM scaffold would improve early engraftment and support the survival of grafted cells and functional recovery post-transplantation.

**Keywords:** PuraMatrix hydrogel, Human stem cells, Traumatic brain injury

**\*Corresponding Author:** Ali Gorji

**Email:** gorjial@uni-muenster.de