



# The 1<sup>st</sup> International Neuroinflammation Congress and 1<sup>st</sup> Student Festival of Neuroscience

Shefa Neuroscience Research Center, Tehran, Iran, 11-13 April, 2017

*The Neuroscience Journal of Shefaye Khatam*

Volume 5, No. 2, Suppl 2

## Oral Presentation

### HTLV-1 and Inflammation

Fariba Zemorshedi\*

Mashhad University of Medical Sciences, Mashhad, Iran

**Published: 11 April, 2017**

#### **Abstract**

Human T-lymphotrophic virus type 1 (HTLV-1) is a retrovirus that is thought to infect approximately 10-20 million people worldwide. The majority of infected individuals remain asymptomatic carriers lifelong. HTLV-1 is thought to cause both local and systemic inflammatory diseases. The most common inflammatory condition associated with HTLV-1 is HTLV-1 associated myelopathy/Tropical spastic paraparesis (HAM/TSP), which occur in 0.25% to 5% of infected individuals. Migration of HTLV-1 infected lymphocytes from periphery into CNS induce a strong immune response and subsequent development of HAM/TSP, which characterized by progressive spastic paraparesis, bladder dysfunction and mild sensory disturbance. HTLV-1 is also associated with other inflammatory conditions such as myositis, uveitis, dermatitis, Alveolitis/bronchiectasis, arthritis and nephritis.

**Keyword:** Inflammation, HTLV, CNS

**\*Corresponding Author:** Fariba Zemorshedi

**Email:** fz6482@yahoo.com