

# The 1<sup>st</sup> International Neuroinflammation Congress and 1<sup>st</sup> Student Festival of Neuroscience



Shefa Neuroscience Research Center, Tehran, Iran, 11-13 April, 2017

*The Neuroscience Journal of Shefaye Khatam*

Volume 5, No. 2, Suppl 2

## Poster Presentation

### Effect of Hesperetin During Pregnancy and Lactation on Locomotor Activity and Anxiety-Like Behaviors in Animal Model of Autism

Rashin Khalaj<sup>1\*</sup>, Akbar Hajizadeh Moghaddam<sup>1</sup>, Mahbobe Zare<sup>2</sup>, Sedigheh Khanjani Jelodar<sup>3</sup>

<sup>1</sup>Department of Biology, Faculty of Basic Sciences, University of Mazandaran, Babolsar, Iran

<sup>2</sup>Faculty of Herbs, Amol University of Special Modern Technologies, Amol, Iran

<sup>3</sup>Faculty of Biology, Shahid Beheshti University, Tehran, Iran

**Published: 11 April, 2017**

#### **Abstract**

**Introduction:** Autism spectrum disorders (ASD) are a heterogeneous group of neurodevelopmental disorders that are defined with a wide range of behavioral impairments including social and communication deficits. Apart from these core symptoms, a significant number of ASD individuals display higher levels of anxiety. The present study is designed to understand the antioxidant efficacy of hesperetin (HET) on locomotor activity and anxiety-like behaviors in an animal model of ASD. **Materials and Methods:** In the experimental research, pregnant rats were divided into four groups including: control, disease group (injected 500 mg/kg valproic acid at gestational day 13) and treatment groups (received 10 and 20 mg/kg/day hesperetin for 7 weeks during pregnancy and lactation). Locomotor activity and anxiety-like behavioral in offspring were measured in the open-field test. **Results:** The results showed that injection of valproic acid increases locomotor activity and anxiety-like behaviors ( $p \leq 0.001$ ) and treatment of hesperetin effectively decrease locomotor activity ( $p < 0.001$ ) and increases duration that spends in middle area of Open field ( $p < 0.01$ ) compared with disease group. **Conclusion:** The results showed that oral administration of hesperetin improves locomotor disorder and anxiety-like behaviors in valproic acid model of autism-like.

**Keywords:** Locomotor Activity, Anxiety, Autism, Valproic acid

**\*Corresponding Author:** Rashin Khalaj

**Email:** r.khalaj70@gmail.com