

# The 1<sup>st</sup> International Neuroinflammation Congress and 1<sup>st</sup> Student Festival of Neuroscience



Shefa Neuroscience Research Center, Tehran, Iran, 11-13 April, 2017

*The Neuroscience Journal of Shefaye Khatam*

Volume 5, No. 2, Suppl 2

## Poster Presentation

### Stem Cells as Neuroinflammatory Modulator in TBI: A Narrative Review

Mohsen Saberifar<sup>1</sup>, Mohammad Amin Bayat Tork<sup>1</sup>, Sajjad Sahab Negah<sup>2\*</sup>

<sup>1</sup>Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>2</sup>Neuroscience Department, Mashhad University of Medical Sciences, Mashhad, Iran

**Published: 11 April, 2017**

#### **Abstract**

Traumatic brain injury (TBI) is physical damage to the brain structure which has a high global rate of mortality and morbidity. TBI can cause intense inflammatory response due to accumulation of leukocytes in cerebral matrix and activation of microglia. Microglia can differentiate into M1 macrophages or M2 macrophages following the changes in biochemical properties of brain tissue. M1 sub type release cytotoxic substances that are toxic to neurons but M2 cells are anti-inflammatory neuroprotective subtype. As the time passes after TBI, the amount of M1 cells begin to increase and fraction of M1:M2 rises. Results show that use of Stem Cells can modulate inflammatory responses of immune system. Transplantation of stem cells into injury site increases M2/M1 ratio as a result of inducing M1 macrophages apoptosis. Different types of stem cells have different mechanisms for anti-inflammatory responses. Even exosomes derived from stem cell can affect the functional recovery and reduce neuroinflammation after TBI. Human Mesenchymal stem cells (hMSCs) are most used in TBI cases due to their immunomodulatory impact and therapeutic effects on recovery of motor and cognitive function. Although many studies conducted to determine effects of hMSCs on TBI prognosis, further investigations are required to support clinical use of hMSCs. Specific attention should be paid to role of growth factors and motifs in suppressing inflammatory responses. Future studies are needed to determine the efficacy of combined therapy.

**Keywords:** Neuroinflammation, TBI, Stem Cell, hMSC, Bioactive motif

**\*Corresponding Author:** Sajjad Sahab Negah

**E-mail:** sahabsajad@yahoo.com