

The 2nd International Neuroinflammation Congress and 2nd Student Festival of Neuroscience



Shefa Neuroscience Research Center, Tehran, Iran, 17-19 April, 2018

The Neuroscience Journal of Shefaye Khatam

Volume 6, No. 2, Suppl 1

Poster Presentation

The Relationship between Spinal Cord Injury and Neuroinflammation and Treatment Methods

Atiyeh Rezaei^{1*}, Parisa Memari Fard¹, Tahereh Naji²

¹Pharmaceutical Science Branch, Islamic Azad University, Tehran, Iran

²Department of Basic Science, Pharmaceutical Branch, Islamic Azad University, Tehran, Iran

Published: 17 April, 2018

Abstract

Spinal cord injury (SCI) is usually caused by a physical factor, especially like burst fracture. Its primary phase involves displacement and physical accidents for the spinal cord, which have two factors of depth and speed of impact. In this phase, most damaged cells are oligodendrocytes in white matter. The secondary phase involves a cascade of cellular and molecular events that progresses rapidly and can cause neuroinflammation. In this phase, The leukocytes and microglia that are in injured region, accelerating the development of neuroinflammation by making unknown species of oxygen reactions. Our goal is to investigate the factors that reduce or improve neuroinflammation. Immunoglobulin G can decrease the activity of leukocytes and microglia and rate of neuroinflammation. Also, the use of useful antioxidants can be effective in reducing neuroinflammation. It seems that presence of immunoglobulin G with an antioxidant can reduce the amount of neuroinflammation caused by SCI seriously. For example, research has shown that combination of palmitoylethanolamid (an endogenous lipid-neuroprotection) and luteolin (antioxidant) can reduce neuroinflammation. also, Ethanol is extracted from black chokeberry (*Aronia melanocarpa* L.) has anti-inflammatory effects. Other studies have also shown that transplantation of bone marrow mesenchymal stem cells (BMCs) and cerebral dopamine neurotrophic factor can inhibit the process of neuroinflammation and decrease production of proinflammation. in addition, it can be effective in regenerating and repairing damaged neurons in SCI. Perhaps a combination of stem cell transplantation and antioxidant effects can help to prevent the development of neuroinflammation and regeneration damaged neurons.

Keywords: Spinal Cord Injury, Neuroinflammation

***Corresponding Author:** Atiyeh Rezaei

Email: rezaiai@yeh@gmail.com