



The 2nd International Neuroinflammation Congress and 2nd Student Festival of Neuroscience

Shefa Neuroscience Research Center, Tehran, Iran, 17-19 April, 2018

The Neuroscience Journal of Shefaye Khatam

Volume 6, No. 2, Suppl 1

Poster Presentation

Agmatine Protects Against Intracerebroventricular Streptozotocin-Induced Water Maze Memory Deficit, Hippocampal Apoptosis and Akt/GSK3 β Signaling Disruption

Maryam Moosavi¹, Amir Hossein Zarifkar^{1, 2*}, Yaghoub Farbood², Mahin Dianat², Alireza Sarkaki², Rasoul Ghasemi³

¹Department of Physiology, Shiraz University of Medical Sciences, Shiraz, Iran

²Department of Physiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences (AJUMS), Ahvaz, Iran

³Department of Physiology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Published: 17 April, 2018

Abstract

Intracerebroventricular streptozotocin (STZ) treatment has been described as a suitable model for sporadic Alzheimer's disease (sAD). Centrally administered STZ decreases insulin and insulin receptors in the brain and interrupts PI3/Akt signaling pathway and GSK-3 β . Additionally it raises Bax/Bcl-2 ratio and prompts hippocampal apoptosis. Agmatine, a polyamine derived from L-arginine decarboxylation, is recently shown to exert some neuroprotective effects. This study aimed to assess if agmatine reverses STZ-induced memory deficits and Akt/GSK-3 β signaling disruption and apoptosis in the hippocampus. Adult male Sprague-Dawley rats weighing 200-250 g were used in this study. The canules were implanted bilaterally into lateral ventricle. STZ was administered on days 1 and 3 (3 mg/kg). Agmatine treatment (40 or 80 mg/kg) was started from day 4 in an every other day manner and continued till day 14. The animal's learning and memory capability was assessed on days 15-18 using Morris water maze. After complement of the behavioral studies the hippocampi was isolated and the amounts of hippocampal cleaved caspase 3 (the landmark of apoptosis), Bax/Bcl-2 ratio, total and phosphorylated forms of GSK-3 β and p-Akt were analyzed by western blot. The results showed that agmatine in 80 but not 40 mg/kg reversed the memory loss induced by STZ. Western blot analysis revealed that STZ induced elevation of caspase-3; Bax/Bcl-2 ratio and disrupted Akt/GSK-3 β signaling in the hippocampus. Agmatine prevented apoptosis and Akt/GSK-3 β signaling alteration induced by STZ. This study disclosed that agmatine treatment avert not only STZ-induced memory deterioration but also hippocampal apoptosis and Akt/GSK-3 β signaling interruption.

Keywords: STZ, Agmatine, Learning and Memory, Apoptosis, Akt, GSK-3 β

***Corresponding Author:** Amir Hossein Zarifkar

E-mail: amirhosseinzarifkar13@yahoo.com