ين رف

The 2nd International Neuroinflammation Congress and 2nd Student Festival of Neurosience

Shefa Neuroscience Research Center, Tehran, Iran, 17-19 April, 2018

The Neuroscience Journal of Shefaye Khatam

Volume 6, No. 2, Suppl 1

Poster Presentation

Depression and the Relation of These Symptoms with Plasma Cortisol Level in Individuals Dependent and Independent to Methamphetamine, a Cross-Sectional Study

Bijan Pirnia^{1, 2*}, Kambiz Pirnia³

¹Department of Psychology, Faculty of Humanities, University of Science and Culture, Tehran, Iran ²Behavioral Sciences Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran ³Technical Assistant in Bijan Center for Substance Abuse Treatment, Tehran, Iran

Published: 17 April, 2018

Abstract

Methamphetamine is a psychostimulant drug that is highly addictive and causes epigenetic changes that is associated with symptoms of depression. The aim of the present study is comparing age and gender differences in individuals dependent and independent to methamphetamine and examining the relation of depression with the level of cortisol. In a cross-sectional study, 55 methamphetamine users with diagnosis of depression (29 men and 26 women) and also 65 non-users depressed patients (30 men and 35 women) among who referred to three stimulant treatment centers in Tehran were selected using purposeful sampling method and were assigned into four groups. Evaluating the level of plasma cortisol hormone was done using radioimmunoassay method (RIA) and depression symptoms were evaluated using Depression Questionnaire. Data were analyzed using chi-square test, multivariate variance analysis and Pierson correlation. The results showed the level of cortisol in two groups of male and female Meth users was higher than two groups of non-users (P<0.05). Also, there is a significant relationship between the level of cortisol and depression in Meth users (P<0.05). This relationship is stronger in women than men. Also, age index in users had lower than two groups of non-users (P<0.05). The findings of the present study can be useful in the process of preventing and treating addiction. Using chemotherapy in creating changes in cortisol levels with the aim of controlling usage relapse can be an appropriate path for future researches in this field.

Keywords: Depression, Plasma Cortisol Level, Methamphetamine

*Corresponding Author: Bijan Pirnia

E-mail: bijanpirnia@yahoo.com

