ثفارغ

The 2nd International Neurotrauma Congress & the 4th International Roads Safety Congress

Shefa Neuroscience Research Center, Tehran, Iran, 18-20 February, 2015

The Neuroscience Journal of Shefaye Khatam

Volume 2, No. 4, Suppl. 3

Workshop Presentation

Animal Models of Traumatic Brain Injury

Maryam Safahani^{1, 2*}, Hadi Aligholi^{1, 3}

¹Shefa Neuroscience Research Center, Khatam Alanbia Hospital, Tehran, Iran.

²Department of Nutrition, Tehran University of Medical Sciences, Tehran, Iran.

³Department of Neuroscience, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran.

Published: 18 February, 2015

Abstract

Traumatic brain injury (TBI) leads to tissue damage by primary and secondary mechanisms. Several factors such as location, nature and severity of the primary injury, age, health, sex, medication, alcohol and drug use, and genetics influence the pathophysiology of TBI in clinic, so TBI is a heterogeneous event and consequently, TBI modeling faces some difficulties. In the fluid percussion injury model, a fluid pressure pulse is generated to the intact dura through a craniotomy and produces a combination of focal and diffuse neuronal injury. The controlled cortical impact injury model uses a pneumatic or electromagnetic impact device to drive a rigid impactor onto the intact dura and creates tissue loss, haematoma, axonal injury and concussion. In the penetrating ballistic-like brain injury, projectiles are transmitted with high energy to produce a cavity in a defined area of the brain. In the weight-drop model, a weight falls from specified height. In the blast brain injury model, trauma can be caused by the primary injury related to the blast. By choosing a proper animal model, we can address the biomechanical aspects of brain injury and assess the potential treatments.

Keywords: Traumatic Brain Injury, Animal Models.

*Corresponding Author: Maryam Safahani

E-mail: maryamsafahani@yahoo.com