

The 2nd International Neurotrauma Congress & the 4th International Roads Safety Congress



Shefa Neuroscience Research Center, Tehran, Iran, 18-20 February, 2015

The Neuroscience Journal of Shefaye Khatam

Volume 2, No. 4, Suppl. 3

Poster Presentation

The Neuroprotective Roles of NMDA Receptors Antagonist Against with Lesion in Central Nucleus of Amygdala

Milad Ahmadi^{1,2*}, Azadeh Sajadian¹, Hadi Aligholi^{1,3}, Babak Khodaie¹, Ahmad Ali Lotfinia¹, Mahmoud Lotfinia^{1,4}

¹Shefa Neuroscience Research Center, Khatam Alanbia Hospital, Tehran, Iran.

²Faculty of Veterinary Medicine, Islamic Azad University, Karaj Branch, Karaj, Iran.

³Department of Neurosciences, School of Advanced Technologies in medicine, Tehran University of Medical Sciences, Tehran, Iran.

⁴Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Published: 18 February, 2015

Abstract

The N-Methyl-D-aspartic acid (NMDA) receptor is one of the specific types of ionotropic receptors which lie to glutaminergic system. It has been widely accepted that NMDA receptor neurons promote anxiety, in humans as well as in animal models. However, in the previous study seems inhibiting of this receptors decreased the protective role in neural cell. Demonstration of NMDA receptor activity had neurodegeneration effect. The hypothesis of this study based on evaluated neuroprotective effect of NMDA receptor inhibition in the central nucleus of the amygdala (CeA) which have role in anxiety. In this study, we confirm that NMDA antagonist (MK801) is neuroprotective when CeA had confronted regional electrical lesion. Kindling of amygdala by electrical lesion which presented by stimulation of the CeA area after microinjection of MK801 (0.5 and 0.75 μ g/kg) in the kindling area is may indicate neuroprotection from secondary injury. It seems the rats were confronted electrical lesion indicated specific decreases in the percentage of open arm time and percentage of open arm entries in elevated plus maze. However, in histopathological study, the mean of dark neurons of CeA in the MK801 group was less than lesion group. We concluded that inhibited of NMDA receptors in CeA neurons may contribute to induced neuroprotection by electrical lesion.

Keywords: NMDA receptor, Neurodegenerative, Dark Neuron.

***Corresponding Author:** Milad Ahmadi

E-mail: pmiladz@gmail.com