# The Second International Anxiety Congress



Shefa Neuroscience Research Center, Tehran, Iran, 1-3 October, 2014

# The Neuroscience Journal of Shefaye Khatam

Volume 2, No.3, Suppl 1

## Poster Presentation

Research

### **Anxiety and Depression among Hospitalised Patients in Mashhad**

Azade Fazli-Shahri1\*, Parisa Samadi1, Roya Samadi2, Masoud Khoshraftar Roodi1, Mehri Baghban Haghighi3

<sup>1</sup> Neuroscience Research Group, Faculty of Medicine, Islamic Azad University Mashad Branch, Mashad, Iran.

<sup>2</sup> Psychiatry and Behavioral Science Research Center, Mashad University of Medical Sciences, Mashad, Iran. <sup>3</sup> General Physician, Mashad, Iran.

Published: 1 Oct 2014

#### **Abstract**

Nonpsychiatric hospitalized patients are frequently exposed to psychiatric disorders, such as anxiety and depression, which significantly influence the patient's subsequent admission days and the rate of recovery. The aim of this study was to evaluate the prevalence of anxiety and depression among hospitalised patients in Mashhad hospitals and to compare the level of anxiety in patients admitted to different departments. This study assessed 264 patients (166 women and 98 men) admitted to internal medicine, general surgery, obestetrics-gynecology and neurology departments of four hospitals in Mashhad during 2012. The level of anxiety and depression was evaluated by hospital anxietydepression scale(HADS). Demographic characteristics and information from medical records was extracted. Data analysis showed that anxiety is related to gender (P<0.001), economic status (P=0.004), employment status (P<0.001), history of chronic medical illness (P=0.004), history of psychiatric illness (P<0.001) and history of usage of psychiatric medicines (P < 0.001). Depression was related to age (P < 0.001, positive correlation), level of education (P = 0.002, illiterate>secondary and tertiary education), economic status (P<0.001, poor>moderate), employment status (P=0.011, students>others), history of chronic medical illness (P<0.001), history of psychiatric illness (P<0.001) and history of usage of psychiatric medicines (P < 0.001). High prevalence of anxiety and depression among hospitalised patients, mainly in women, reminds us the necessity of performing psychiatric consults in physical patients in order to decrease the burden of disease, hospitalisation days and omission of coexisted psychiatric disorders.

**Keywords:** Anxiety, Depression, Hospitalized Patients, Demography, Mashhad.

\*Corresponding Author: Azadeh Fazli-Shahri

Email: azadehfazli@yahoo.com