

The 2nd International Neurotrauma Congress & the 4th International Roads Safety Congress

Shefa Neuroscience Research Center, Tehran, Iran, 18-20 February, 2015

The Neuroscience Journal of Shefaye Khatam

Volume 2, No. 4, Suppl. 3

Poster Presentation

Minimally Conscious State Improvements with Thalamic Ventralis Intermedius Nucleus Stimulation after Severe Traumatic Brain Injury

Milad Ahmadi*, Babak Khodaie, Ahmad Ali Lotfinia, Mahmoud Lotfinia

Shefa Neuroscience Research Center, Khatam Alanbia Hospital, Tehran, Iran.

Published: 18 February, 2015

Abstract

Methods of modulating brain neural plasticity, improving memory and learning, improving recovery from traumatic brain injury, preventing epilepsy, treating memory disorders, and treating persistent impairment of consciousness in humans and animals by nerve stimulation are provided. Widespread loss of cerebral connectivity is assumed to underlie the failure of brain mechanisms that support communication and goal-directed behavior following severe traumatic brain injury. This study was tried to clarify with the effect of deep brain stimulation in patients who had severe traumatic brain injury. Recent studies have shown unexpected preservation of large-scale cerebral networks in patients in the minimally conscious state (MCS), a condition that is characterized by intermittent evidence of awareness of self or the environment. These findings indicate that there might be residual functional capacity in some patients that could be supported by therapeutic interventions. This study was followed after 6-mounth indicated that bilateral deep brain electrical stimulation (DBS) of the ventral is intermedius thalamus modulates behavioral responsiveness in a patient who remained in MSC for 6 years following traumatic brain injury before the intervention. Primary outcome measures was mediated by the frequency of specific cognitively behaviors and functional limb control and oral feeding (secondary outcome measures) increased during periods in which DBS was on as compared with periods in which it was off. Our studies demonstrated early DBS treatment discontinuation for patients with only inconsistent interactive behaviors and motivate further research to develop therapeutic interventions.

Keywords: Traumatic Brain Injury, Minimally Conscious State, Deep Brain Stimulation, Neural Plasticity.

*Corresponding Author: Milad Ahmadi

E-mail: dr.miladahmady@gmail.com